<u>Cite this as:</u> Zi-li MA, Yang GAO, Hai-tian MA, Liu-hai ZHENG, Bin DAI, Jin-feng MIAO, Yuan-shu ZHANG, 2016. Effects of taurine and housing density on renal function in laying hens *Journal of Zhejiang University-Science B (Biomedicine & Biotechnology)*, 17(12): 952-964.

http://dx.doi.org/10.1631/jzus.B1600014

Effects of taurine and housing density on renal function in laying hens

Key words: Rearing pattern, Welfare, Taurine, Laying hens, Renal impairment

Research Summary

This study mainly focused the possibility of ameliorating renal impairment, increasing productivity, enhancing health and promoting welfare in laying hens by reducing housing density and by dietary supplementation with taurine and summarized the key roles they played in the following aspects:

- Histological examination
- Quantification of SCr, BUN, SUA, NAG and GLU
- Changes in iNOS
- Effect of taurine and different housing modes on NF-κB DNA binding activity


- Real-time quantitative RT-PCR analysis of gene expression
- The detection of TNF-α levels by ELISA
- Activities of SOD and levels of T-AOC and MDA in the kidney

Innovation points

- Introduction of the putative protective effects of supplemental taurine and reduced housing density on renal function in laying hens.
- Summary of taurine supplementation, and rearing free-range or in low-density housing can lessen morphological renal damage, inflammatory mediator levels and oxidation levels and increase anti-oxidation levels.
- Emphasis of taurine supplementation and a reduction in housing density can ameliorate renal impairment, increase productivity, enhance health and promote welfare in laying hens







Innovation points

- Table 1 Ingredient composition and nutrient content of diets
- Table 2 Primer sequences of targeted genes and β-actin
- Fig. 1 The clinical photos of laying hens
- Fig. 2 Histology of the kidney in laying hens. Renal sections were stained with H&E (original magnification 200 X)
- Fig. 3 SCr, BUN, SUA, NAG and GLU levels
- Fig. 4 Changes in SOD activity and T-AOC, MDA levels in kidney
- Fig. 5 Effect of taurine on NF-kB DNA binding activity by EMSA analysis in renal tissue from laying hens
- Fig. 6 Changes in IFN- γ , IL-4, IL-10 and PPAR- α gene expression in laying hen kidneys
- Fig. 7 Changes in iNOs, TNF- α levels and TNF- α gene expression in the kidneys of laying hen