



Review

<https://doi.org/10.1631/jzus.B2300776>



Structure of myelin in the central nervous system and another possible driving force for its formation—myelin compaction

Qi SHAO¹, Simin CHEN¹, Tian XU², Yuyu SHI¹, Zijin SUN¹, Qingguo WANG¹,
Xueqian WANG^{1✉}, Fafeng CHENG^{1✉}

¹College of Traditional Chinese Medicine, Beijing University of Chinese Medicine, Beijing 100029, China

²Department of Gastroenterology, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences, Beijing 100730, China

Abstract: Myelin formation is considered the last true “invention” in the evolution of vertebrate nervous system cell structure. The rapid jumping pulse propagation achieved by myelin enables the high conduction speed that is the basis of human movement, sensation, and cognitive function. As a key structure in the brain, white matter is the gathering place of myelin. However, with age, white matter-associated functions become abnormal and a large number of myelin sheaths undergo degenerative changes, causing serious neurological and cognitive disorders. Despite the extensive time and effort invested in exploring myelination and its functions, numerous unresolved issues and challenges persist. In-depth exploration of the functional role of myelin may bring new inspiration for the treatment of central nervous system (CNS) diseases and even mental illnesses. In this study, we conducted a comprehensive examination of the structure and key molecules of the myelin in the CNS, delving into its formation process. Specifically, we propose a new hypothesis regarding the source of power for myelin expansion in which membrane compaction may serve as a driving force for myelin extension. The implications of this hypothesis could provide valuable insights into the pathophysiology of diseases involving myelin malfunction and open new avenues for therapeutic intervention in myelin-related disorders.

Key words: Myelin; Central nervous system; White matter; Myelin compaction

1 Introduction

Myelin is a multilayered sheath that envelops the outside of nerve cell axons. In the central nervous system (CNS) of mammals from marsupials to humans, myelin sheaths are derived from oligodendrocytes (OLs), which are derived from morphologically complex precursor cells (oligodendrocyte precursor cells (OPCs) or neural/glia antigen 2 (NG2) cells). OPCs are a type of glial cell that exists in the CNS and originates from neural stem cells (NSCs). They are the largest dividing population of neural cells, distributed evenly and accounting for an average of 5% to 10% of all CNS cells (Tepavčević and Lubetzki, 2022).

Mature OLs create myelin by wrapping their own cell membranes around axons in a spiral shape, ultimately becoming multilayered sheaths that cover long axon segments. The fundamental function of myelin is saltatory conduction from one node of Ranvier to the next, thereby increasing nerve impulse conduction velocity (Zalc et al., 2008). High conduction speed is the basis of motor, sensory, and cognitive functions. Therefore, the formation of myelin is regarded as the last true “invention” in vertebrate evolution in the structure of nervous system cells (Nave, 2010).

White matter is the primary component of the CNS in humans (accounting for 60% of brain volume) and is composed of a large number of myelinated nerve fibers (Krafft et al., 2012). In contrast, gray matter is composed of a large number of nerve cell bodies and synapses, and contains less myelin than white matter. Owing to its location at the end of arterial circulation, white matter is particularly susceptible to reduced blood flow and oxygenation. Consequently, with age, a significant number of myelin sheaths undergo

✉ Xueqian WANG, wxqbum@126.com

Fafeng CHENG, fafengcheng@gmail.com

Xueqian WANG, <https://orcid.org/0000-0002-7682-3877>

Fafeng CHENG, <https://orcid.org/0000-0002-6742-0742>

Received Oct. 26, 2023; Revision accepted Feb. 22, 2024;

Crosschecked Apr. 22, 2025

© Zhejiang University Press 2025

degenerative changes that lead to severe neurological and cognitive impairments (Liu et al., 2017). Since OLs are primarily responsible for synthesizing myelin, changes in OLs and their precursor cells can significantly affect the structure and function of myelin. Nevertheless, the degeneration of OLs and their precursor cells is exacerbated with age, which may be associated with microglia- and astrocyte-mediated inflammation, mitochondrial dysfunction, and iron accumulation (Correale and Ysraelit, 2022). In this review, we elucidate the structure and function of myelin and describe the process of myelin formation to lay a foundation for a better understanding of myelin-related diseases. Finally, we introduce the epidemiology of myelin-related diseases and describe their potential mechanisms of action.

2 Structure and function of myelin

2.1 Major dense line

Alternating major dense lines and intraperiod lines are the two characteristic periodic morphological

features of the myelin sheath. The major dense lines, 3 nm in width, are formed by the tightly packed intracellular (cytoplasmic) surfaces located between the inner membranes of the lipid bilayers (Aggarwal et al., 2011; Kister and Kister, 2023). This unique structure is stabilized by multiple adhesive mechanisms, in which myelin basic protein (MBP) is essential. MBP is the second most abundant protein in the CNS myelin, accounting for 30% of the mass of myelin protein in the CNS. MBP tightly binds the two negatively charged cytoplasmic leaflets of the myelin membrane that form the major dense lines, and is referred to as an “executive” molecule of the myelin membrane, playing a key role in the formation of compact myelin (Moscarello, 1997) (Fig. 1a). MBP is essentially an intrinsically disordered protein; however, upon binding to membranes, MBP adopts α -helical and β -sheet structures (Harauz et al., 2004) (Fig. 1b). Membrane binding alters the properties of MBP, promoting self-interaction into a densely packed protein phase that occupies the major dense lines and tightly binds the cytoplasmic surface of the bilayer together to limit the entry of most proteins into the myelin sheaths




Fig. 1 Schematic diagram of the myelin structure in the central nervous system (CNS). (a) Longitudinal section of myelin; (b) Intracellular lipid bilayers and cell membranes; (c) Cross section of myelin, where c-1 is a node, c-2 a paranode, c-3 a juxtapanode, c-4 an internode, and c-5 a gap junction in an oligodendrocyte; (d) Gap junction in oligodendrocyte and astrocyte networks. Caspr: contactin-associated protein; CNP: 2',3'-cyclic nucleotide 3'-phosphodiesterase; Cntn: contactin; Cx: connexin; F-Actin: filamentous actin; IPL: intraperiod line; K⁺Ch: K⁺ channel; MAG: myelin-associated glycoprotein; MDL: medulla; MBP: myelin basic protein; Na⁺Ch: Na⁺ channel; NECL4/1: nectin-like 4/1; NF186/155: 186/155-kDa isoform of neurofascin; PLP: proteolipid protein.

(Raasakka et al., 2017; Snaidero et al., 2017). Force-distance measurements indicate that maximal adhesion force and minimal cytoplasmic spacing occur when each negative lipid in the membrane can bind to a positively charged arginine or lysine residue on MBP (Min et al., 2009). In addition to its structural importance (through interactions with lipids and myelin-associated proteins), MBP may play a regulatory role in myelination because it participates in pathways that promote both neuron survival and death, such as binding to MAGEB2/D2 proteins associated with neurotrophin receptor p75NTR1 (Smirnova et al., 2021).

Both an excess and a deficiency of MBP can cause the repulsion of myelin bilayers (this pertains only to the cytoplasmic sides of myelin bilayers) and may lead to the destruction of myelin (demyelination). Elevated MBP concentrations induce excessive positive charges that generate both interlayer electrostatic repulsion and pathological water gap expansion, resembling demyelination *in vivo*. A phase transition is observed across highly swollen bilayers, indicating the formation of a weak, dilute, yet extensive gel-like structure of MBP, which allows it to collapse under a compressive pressure of approximately 1 MPa. When the swelling pressures are not too severe, the bilayers can be pressed together again to almost the same separation, within 0–2 nm, even from distances as large as 80 nm (Min et al., 2009). Shiverer mice, which lack expression of MBP, form less myelin, which is less stable than that in wild-type mice. The myelin formed by shiverer mice consists of loosely compacted layers and is at most four layers thick. Although some basic structural elements of the myelin remain, such as radial components (RCs) and axon–glial connections, most of the axonal surface is completely devoid of myelin (Brady et al., 1999; Snaidero et al., 2014). In contrast to mice lacking basic components of paranodal axo-glial cell junctions (such as contactin-associated protein (Caspr), contactin-1, and 155-kDa isoform of neurofascin (NF155)), shiverer mice do not exhibit severe axonal degeneration (Sherman et al., 2005; Pillai et al., 2009). This may be because non-myelinated nerve fibers in shiverer mice are still able to propagate action potentials, albeit in an immature continuous manner, whereas mice lacking paranodal structures suffer from conduction block caused by current leakage (McNamara et al., 2023).

2.2 Intraperiod line

An intraperiod line is formed by the outer surface of the myelin membrane lipid bilayer. The intraperiod line is wider than the main dense lines when observed using electron and X-ray microscopy (Karthigasan and Kirschner, 1988; Kirschner et al., 1989; Snaidero et al., 2014). According to most scholars, the intraperiod line measures 4–5 nm, whereas the main dense line is about 3 nm wide (Raasakka et al., 2019; Kister and Kister, 2023). The values calculated by Min et al. (2009) based on the cable theory equation set, align closely with these measurements (Figs. 1a and 1b). Proteolipid protein (PLP) is the most abundant myelin protein in the CNS, accounting for 38% of the total myelin protein mass. PLP is a highly conserved hydrophobic protein composed of 276 amino acids, contains four transmembrane segments and two disulfide bonds, and covalently binds with lipids (at least six palmitic acid groups in mammals) (Weimbs and Stoffel, 1992). PLP exists in two isoforms, a larger isoform weighing 30 kDa and a shorter isoform (DM20) weighing 26 kDa, which lacks a specific sequence of 35 amino acids (Ruskamo et al., 2022). PLP and DM20 have strong lipophilicity and are expected to contain fatty acid portions connected by several cysteines. PLP interacts with the membrane and has a high affinity for CH-rich lipid rafts. PLP and DM20 play important roles in forming and maintaining the axon integrity of myelin periodic lines. One key function of PLP is to bind myelin lamellae together and enhance the compaction of myelin lipids, thereby increasing the physical stability of myelin and supporting axon-myelin metabolism (Ruskamo et al., 2022). Therefore, a high level of PLP is required in the myelin to maintain its integrity. A 50% decrease in PLP content can lead to ultrastructural and axonal pathological changes in myelin, including unfolding of myelin, splitting of lamellae, and the formation of axonal spheroids (Lüders et al., 2019).

The adhesion function of PLP has been a topic of controversy. In Pelizaeus-Merzbacher patients, a more severe clinical phenotype results from mutated PLP than from an absence of PLP (Stadelmann et al., 2019). Several studies have shown that PLP-deficient mice exhibit moderate phenotypic changes, including slightly reduced or delayed formation of myelin with small-diameter axons, an increased number of cytoplasmic channels in compact myelin, insufficiently

compacted myelin, and axonal damage (Ruskamo et al., 2022). The lack of PLP also damages the abundance of cholesterol in myelin, axonal integrity, and motor ability (Lüders et al., 2019). However, in mice whose PLP was completely knocked out, these severe phenotypes have not been confirmed. Instead, compact myelin sheaths, which are even more tightly compressed and periodically reduced, have been demonstrated (Klugmann et al., 1997). Recent studies have found that PLP does indeed have adhesion; however, even without PLP, the extracellular outer surfaces of the myelin layer membrane will stick together. Therefore, it is not the expression of PLP but rather the loss of repulsion between adjacent membrane bilayers that may drive the compaction of myelin at the site of PLP (Bakhti et al., 2013). All these results suggest that while PLP is not necessary for myelin formation, it is necessary for axonal integrity. Furthermore, when PLP is overexpressed in transgenic mice, the CNS exhibits poor myelin development and demyelination, and disease severity is correlated with PLP expression levels (Kagawa et al., 1994). It is unclear why abnormal levels of PLP lead to cell death; however, changes in the proportion of myelin membrane components may lead to the accumulation of toxic substances inside cells (Stadelmann et al., 2019).

2.3 Non-compact myelin

When MBP binds to the surface of two adjacent cytoplasmic membranes, it drives the membrane zipper on the cytoplasmic surface of the myelin bilayer. This process causes cytoplasmic solute extrusion and forms a narrow and dense protein phase. To provide directionality during compaction and exclude the formation of isolated non-compacted cytoplasmic pockets in the myelin, there must be a mechanism to counteract MBP-mediated membrane compaction. 2',3'-Cyclic nucleotide 3'-phosphodiesterase (CNP) appears to be a protein that accomplishes this function (Snaidero et al., 2014, 2017) (Fig. 1b). CNPs are abundant in the CNS, where they make up 4% of the total proteins of the myelin sheath (Gravel et al., 2009). They are located in OL cell bodies and non-compact myelin, i.e., tongue-in and tongue-out protrusions and paranodal loops (Edgar et al., 2009). During development, CNP is highly up-regulated in OLs before myelination and remains constant throughout life, playing a crucial role in the formation and lifelong maintenance of myelin (Lee

et al., 2005). First, CNP can directly bind to the actin cytoskeleton to form a robust structure that can counteract the adhesion force exerted by MBP molecules (Snaidero et al., 2017). Second, as a dual-function protein, both a phosphodiesterase and a membrane-binding microtubule-associated protein (MAP), CNP can act as a linker protein between microtubules and cell membranes to regulate microtubule dynamics (Bifulco et al., 2002). During myelin formation, microtubules provide transport pathways for vesicles and *Mbp* messenger RNA (mRNA) (Müller et al., 2013). Studies have also shown that CNP binds to poly(A)⁺ mRNA in vivo and inhibits in vitro translation in a dose-dependent manner, possibly regulating mRNA expression in OLs of the CNS (Gravel et al., 2009).

In the absence of CNP, myelin compaction proceeds more rapidly (Snaidero et al., 2014). Moreover, CNP-deficient mice exhibit progressive axonal pathology characterized by axonal swelling and spheroid formation (Lappe-Siefke et al., 2003), with a lower number of cytoplasmic channels within their myelin sheaths (Snaidero et al., 2017). When MBP levels are reduced, cytoplasmic channels are restored in CNP-deficient mice and the pathology of large-caliber axons is reduced (Snaidero et al., 2017), indicating that MBP has an antagonistic effect on CNP. Similarly, overexpression of CNP occurs in areas lacking compact myelin (Gravel et al., 1996) and increases the amount of cytoplasm within the myelin. Therefore, CNP plays a crucial role in generating myelin and maintaining axonal physiology.

2.4 Radial components

In the CNS, the adhesion force between adjacent myelin layers converges at the RC. The normal CNS myelin compaction along the outer surface of the lamella is believed to be maintained by the interaction between PLP molecules in contiguous layers and the binding at the junction of the RC. One of the components of these tight connections is claudin-11, which is an abundant myelin protein. The RC consists of multiple layers, with most RC elements located in the outer and inner tongues of the myelin. RC elements must be interconnected in some ways between myelin layers to explain their formation and maintenance in register, even across different myelin sheaths (Rosenbluth et al., 2006). Claudin-11 forms a tight junction barrier that enhances the insulating properties of CNS

myelin, but this enhancement is typically significant only for small axons with thin sheaths and is negligible for large axons (Devaux and Gow, 2008).

Initial research discovered that mice deficient in claudin-11 lacked RCs and exhibited mild tremors, motor defects, and gait and electrophysiological abnormalities. These abnormalities included a 50% reduction in conduction velocity in small-diameter axons (Gow et al., 1999). Subsequent studies confirmed this finding. Devaux and Gow (2008) performed an electrophysiological analysis of claudin-11-deficient mice and reported that the conduction velocity in small-diameter fibers was severely slowed down. However, the length and thickness of the myelin were normal. These abnormalities were not caused by disruption of the myelin structure or distribution of ion channels in axons, but rather by changes in the biophysical properties of myelin. Specifically, changes in the function of the myelin barrier suggest that the main function of claudin-11 is to form a diffusion barrier rather than to provide mechanical stability. In support of this idea, Denninger et al. (2015) found that myelin lacking claudin-11 had stronger permeability to water and small permeants while maintaining the integrity of the myelin structure. However, other studies have shown that when both *claudin-11* and *PLP* genes are knocked out, mice have severe neurological dysfunction, significantly abnormal myelin compaction, and smaller axon diameters. Interestingly, when either of these genes is knocked out, the expression of the other proteins increases. These studies suggest that claudin-11 and PLP serve fundamental structural functions in preserving normal compact myelin and there is redundancy in their functions (Chow et al., 2005). Claudin-11 may have more functions in forming barriers, enhancing the insulation performance of CNS myelin, and maintaining the normal physiological characteristics of axons and relatively less effect on the structure of myelin.

2.5 Gap junctions

Gap junctions (GJs) are channels that directly connect the cytoplasm of coupled cells. They participate in intercellular substance and energy exchange, and information transmission. Their main function is to mediate the transmission of chemical signals between cells and electrical signals across synapses in neurons. GJs allow ions (Ca^{2+} , Na^{+} , and K^{+}), glutamate (Glu), glutathione, cyclic adenosine monophosphate

(cAMP), inositol 1,4,5-triphosphate (IP_3), adenosine triphosphate (ATP), and more to pass through (Lapato and Tiwari-Woodruff, 2018). They play important roles in many aspects of cell metabolism, including proliferation and differentiation, as well as energy transport and maintenance of tissue homeostasis (Fig. 1c). In recent years, GJ intercellular communication (GJIC) channels have been widely studied as emerging membrane protein channels. Almost all differentiated cells in vertebrates communicate through GJs, including red blood cells, smooth muscle cells, epithelial cells, and neurons. Various diseases of the CNS, such as demyelinating diseases, epilepsy, gliomas, and cerebral ischemia-hypoxia injury, have been confirmed to be related to GJIC (Hou et al., 2020). GJ proteins are a multi-gene family-encoded class of membrane proteins consisting of four transmembrane domains connected by two extracellular loops. Six connexin proteins oligomerize to form a hemichannel that allows the diffusion of ions and small molecules through two hemichannels on adjacent cell membranes that form a GJ. In the CNS, OLs express Cx29, Cx32, and Cx47; astrocytes express Cx26, Cx30, and Cx43; microglia express Cx32a, Cx36, and Cx43a; and neurons express Cx30.2, Cx31.1, Cx32, Cx36, Cx40, Cx45, and Cx50 (Lapato and Tiwari-Woodruff, 2018). Communication between glial cells in the CNS is essential for the formation of normal myelin. The absence or disruption of these connections may lead to severe demyelinating diseases and early death.

Cx47 and Cx32 are GJ proteins that have been shown to form functional GJs, such as Cx47/Cx47 and Cx47/Cx43, which promote the diffusion of nutrients, ions, secondary messengers, and small molecules between cells. Cx47 plays a critical role in the formation and repair of myelin in the CNS. On the other hand, Cx32 forms mainly Cx32/Cx32 channels and plays a key role in maintaining the structure and function of myelin, related mainly to the degree of demyelination. The GJs between OLs/astrocytes are composed mainly of Cx47/Cx43 and Cx32/Cx30 channels, with Cx47/Cx43 channels located mainly near the OL cell body and Cx32/Cx30 channels located mainly in the outer layer of the myelin (Maglione et al., 2010) (Fig. 1d). Studies have shown that the absence of Cx47 and Cx32 leads to a slowdown in mouse myelin development, while the simultaneous absence of both leads to severe myelin abnormalities and death within four

months after birth (Tress et al., 2011). Clinically, mutations in the Cx47 promoter lead to Pelizaeus-Merzbacher-like disease (Gotoh et al., 2014), while changes in Cx32 lead to symptoms very similar to those of Charcot-Marie-Tooth disease (Sargiannidou et al., 2015). However, the precise roles of these GJ proteins are still unclear. GJs deserve further in-depth investigation due to their critical potential functions in facilitating metabolic transport, spatial buffering, and electrical coupling between cells.

2.6 Axon–myelin junctions

The connection between the axon and myelin is composed of three main parts: the paranode, juxtaparanode, and internode. During the formation of myelin, glial cells also push axonal proteins to reorganize into specific regions. It is noteworthy that the formation of myelin causes voltage-gated sodium channels to accumulate at the nodes of Ranvier (the gaps between the myelin), which are approximately 1 μm long and correspond to less than 1% of the myelinated axon area (c-1 in Fig. 1). In the CNS, node assembly begins with axoglial junctions formed at paranodes, which act as diffusion barriers (c-2 in Fig. 1). The paranodal junction is the closest and parallel part between the axon and myelin glial cells. As a result of its circumferential wrapping, the outer, uncompacted edge of the glial cell exhibits a series of loops invaginating the axon (Salzer and Zalc, 2016). At paranodes, axonal proteins contactin and Caspr and glial molecule NF155 form a cell adhesion complex. One characteristic of paranodal connections is the regular interval transverse diaphragm emanating from the axon. Paranodal connections act as membrane diffusion barriers, promoting the accumulation of voltage-gated sodium channels at nodes and separating them from potassium channels (Kv1.1 and Kv1.2) in juxtaparanodes (Salzer and Zalc, 2016) (c-3 in Fig. 1).

The loss of paranodal junctions results in the mislocalization of juxtaparanode Kv1 channels to paranodal regions and impairs conduction. In addition to their role as a diffusion barrier, paranodal junctions facilitate nerve conduction by enclosing the myelin on the axon, preventing nodal action currents from being shunted below the myelin, while still leaving a narrow channel connecting the internodal periaxonal space and the perinodal space. In juxtaparanodes near the paranodal region, axonal Caspr2, transient axonal glycoprotein-1 (Tag-1), and glial cell Tag-1 form a

cell adhesion complex that initiates the assembly of Kv1 channel clusters. However, the interaction between Caspr2/Tag-1 and Kv1 channels remains unclear. The special structural domains of these ion channels and cell adhesion molecules (CAMs) are further stabilized by submembrane cytoskeletal complexes in nodes, paranodes, and juxtaparanodes. For more information on this topic, please refer to the studies reported by Susuki et al. (2016), Fehmi et al. (2018), and Pinatel and Faivre-Sarrailh (2021).

Myelin-associated glycoprotein (MAG) plays a crucial role in internodes. It is produced by myelinating glial cells and is located around the entire circumference of the axonal myelin (c-4 in Fig. 1). MAG has a high expression level (1%) in the CNS and exists as two isoforms (large MAG (L-MAG) and small MAG (S-MAG)). L-MAG is produced almost exclusively in the CNS and is the major variant during early development and active myelin formation (Campagnoni, 1988; Trapp, 1990). S-MAG is the major subtype in adult CNS and all age groups of the peripheral nervous system (Harry and Toews, 1998). Although MAG has long been considered an axon growth inhibitor along with Nogo and oligodendrocyte myelin glycoprotein (OMgp) (McKerracher and Rosen, 2015), its effect on axon growth is bidirectional; it promotes axon growth in young neurons and inhibits growth in older neurons, which is an age- and neuron-type-dependent switch. Deletion of MAG leads to accelerated axonal loss in experimental autoimmune encephalomyelitis (an animal model of multiple sclerosis). This suggests that MAG can also promote axon growth and/or protect axons from further degeneration in the adult CNS (Geoffroy and Zheng, 2014). However, further research is needed on the roles of MAG and its receptors in myelin development.

3 Formation and regeneration of myelin

3.1 Maturation of oligodendrocytes

OLs are the sole myelin-forming cells in the CNS and are a major type of glial cell in the brain. They play a crucial role in promoting the propagation of action potentials and supporting the integrity of neurons and axons while providing metabolic and nutritional support for axons. OLs are believed to be derived from OPCs, which differentiate into pre-OLs before finally maturing into myelin OLs (Fig. 2).




Fig. 2 Main markers of different stages of oligodendrocyte development (proliferation, differentiation, and maturation). For a detailed introduction to these biomarkers, please refer to Simons and Nave (2016), Kuhn et al. (2019), Kang and Yao (2022), and Wang et al. (2022). The genes marked in black are markers of oligodendrocyte precursor cell (OPC) proliferation and oligodendrocyte (OL) differentiation and maturation, while those marked in red are inhibitors of OL differentiation or maturation.

OPCs are generated in the ventral forebrain where three distinct groups of OPCs have been identified. These groups are generated in a continuous wave-like pattern with a clear spatiotemporal pattern and migrate from their birthplace to settle in the brain. Although they have different origins, all OPC cell lineages converge at the transcriptional level, which may reflect their contribution to myelin formation. During migration, some neural cells promote transient cell-cell interactions that allow them to play a greater role after integration into the cortical network (Lepienne et al., 2022). OPCs constantly extend and retract during migration and settlement into their final positions. These dynamic and exploratory processes are repulsive to other OPCs, thus determining the uniform density and spacing of these cells (Hughes et al., 2013). After OPCs enter their designated area, some remain in a precursor state while others differentiate into OLs that form myelin.

In rodents, OPCs proliferate and migrate mainly in the postnatal stage, around the second day after birth. Late-stage OPCs account for most OL lineage cells, with a small number of immature OLs also present. Mature OLs appear on the seventh day after birth and initiate CNS myelination. Myelination in most brain regions is almost complete by around 60 d after birth

(Snaidero et al., 2014). In humans, OPCs are first observed in the forebrain at around 10 weeks of gestation. Although OPCs and late-stage OPCs remain the main cell types during this stage, immature OLs are observed between 18 and 28 weeks. Therefore, the OL lineage develops between 18 and 28 weeks, similar to rodents on postnatal Day 2. In human development, MBP-positive OLs are initially observed at around 28 weeks, similar to rodents on postnatal Day 7 (Barateiro et al., 2016).

Mature OPCs undergo different stages of morphological differentiation and express many cell-specific marker proteins and lipids. Although some molecules have opposing functions in OPC proliferation and OL differentiation, the physiological processes they control are not opposite. Transient expression of signaling molecules that drive OPC proliferation and prevent OL differentiation or maturation is necessary for early OPC content control in the CNS, while their continuous transient expression is necessary for myelin formation or remyelination (Wang et al., 2022). The balance between OPC proliferation, OL differentiation and maturation is delicate and complex. Once myelination begins, immature OLs must be properly positioned because mature OLs appear to have little ability to migrate to new locations or delay myelin

synthesis. Once myelination begins, signaling pathways dependent on phosphatidylinositol (3,4,5)-trisphosphate (PtdIns(3,4,5)P₃)/protein kinase B (Akt)/mechanistic target of rapamycin (mTOR) and extracellular signal-regulated kinase 1/2 (ERK1/2) are critical for driving myelination until mature myelin with optimal G-ratios is produced (Simons and Nave, 2016). For more detailed information on the growth, development, and migration of OPCs, please refer to the study by Kang and Yao (2022).

3.2 Recognition and adhesion of axons

During the initial period of myelination, the mechanism by which OLs select axons and their subdomains for myelination remains elusive (Fig. 3). A recent study demonstrated that myelin formation can be initiated by the axon diameter alone without any axonal signals (Mayoral et al., 2018). Other studies have shown that OLs can wrap around artificial glass or polymer matrices and sense the diameter of microfibers, thereby increasing the length of the sheath as the fiber grows (Lee et al., 2012; Bechler et al., 2015). These findings suggest that physical properties such as axon diameter play crucial roles in initial axon selection. However, some axons in the brain between 0.2 and 0.8 μm have myelin while others do not (Waxman and Bennett, 1972), indicating that axon diameter is not the sole determinant of myelin formation. Recent studies have found that blocking action potentials in zebrafish spinal cords leads to incorrect axon targeting and insufficient myelination (Hines et al., 2015). In addition, OLs can myelinate axons (Rosenberg et al., 2008) fixed with paraformaldehyde (PFA) and nanofibers (Lee et al., 2013) in the absence of neural activity. Furthermore, some scholars have found that OLs on the axons of rodents extend or retract according to permissive or repulsive cues, respectively (Thomason et al., 2020). These findings suggest that correct axon targeting is likely to depend on multiple cues, including axon diameter, neural activity, and permissive or repulsive cues on OLs. However, which

factors specifically determine axon selection and how they relate to each other are still unknown.

3.3 Growth of myelin and axon wrapping

Upon finding the target axon, OLs undergo membrane polarization which leads to the accumulation of myelin-specific proteins and lipids in the myelin (Fig. 3). During development, the myelin membranes of OLs can grow up to 5000 μm^2 per day (Pfeiffer et al., 1993). A single OL can myelinate up to 50 axons (de Faria et al., 2021; Pajevic et al., 2023), while mature myelin consists of up to 160 membrane layers (Ruskamo et al., 2022). Several models, such as the “croissant-like” model (Sobotka et al., 2011) and the “yo-yo” model (Pedraza et al., 2009), were initially proposed for the wrapping of myelin. However, with the advent of three-dimensional (3D) electron microscopy techniques and high-pressure freezing techniques for fixed tissues, new models have been proposed for the wrapping of myelin (Kerman et al., 2015). The innermost layer of the myelin is the shortest and extends laterally while the top of the myelin gradually increases with each wrapping. This means that myelin grows through two different but coordinated movements: the leading edge at the inner tongue wraps around the axon (i.e., beneath previously deposited membranes) and extends laterally toward the node. Thus, each myelin layer remains in close contact with the axonal surface and moves forward in a continuous spiral manner toward future nodes where they align and are positioned as paranodal loops. Cytoplasmic channels are present in compact myelin sheaths to provide short connections between the outer and inner tongues that allow newly synthesized membrane components to be transported from cell bodies to inner tongues. Most of these channels close with ongoing development, but residual cytoplasmic channels persist throughout the mature compact myelin sheath (Snaidero et al., 2014; Edgar et al., 2021). The cytoplasmic channels appear to be dynamic and have the potential to reopen after the developmental program is terminated (Snaidero et al., 2014).

3.4 Myelin compaction

Myelin compaction typically occurs after a period of wrapping and simultaneously with wrapping. Although the specific mechanism regulating compaction initiation and corresponding sites is not yet clear,




Fig. 3 Schematic of central nervous system (CNS) myelination.

it is clear that compaction on the cytoplasmic surface starts from the outermost layer of myelin phospholipids and progresses inward. During initial formation, myelin is rich in negatively charged phospholipids (PIP2, PIP3). MBP has a high affinity for negatively charged phospholipids, thereby pulling two bilayers together to play a “zipper” role (Snaidero et al., 2014). However, MBP is synthesized in the distal process (i.e., closest to the axon) (Laursen et al., 2011), while myelin compaction occurs at the proximal end. This phenomenon is similar to PLP because PLP is synthesized on ribosomes in the endoplasmic reticulum and then transported to the myelin membrane through a series of steps (Kippert et al., 2007). One possible explanation is that compaction starts from the site where MBP mRNA is translated locally and then moves inward after binding with other proteins (Laursen et al., 2011). The inner tongue may contain necessary substances produced by MBP, but it cannot provide conditions for compaction because it needs further expansion and extension. This also explains why compaction always starts after several layers of wrapping. To maintain this rate of membrane compaction and exclude the formation of unconsolidated areas, CNP plays an important role. Unconsolidated areas are rich in CNP; therefore, unconsolidated wrapping decreases in the absence of CNP (Snaidero et al., 2014). CNP may have additional functions as myelin sheaths lose some cytoplasmic channels when it is absent (Snaidero et al., 2014), and the amount of cytoplasm in the myelin increases when it is overexpressed (Gravel et al., 1996; Snaidero et al., 2017). This also means that during the growth of the myelin, the rapid extension of the innermost part of the myelin and the accumulation of CNPs between these newly synthesized layers prevent MBP from being compressed prematurely at the front (Snaidero et al., 2014). Under this mechanism, MBP synthesis sites may occur in the innermost tongue, and then MBP spreads to the outer layer where the lamellae are closer together and begin to compact (Stadelmann et al., 2019).

3.5 Node formation

During the lateral extension of myelin sheaths, a special structure called a Ranvier node is formed when two myelin sheaths meet. Myelin phospholipids are electrical insulators. When the membrane at the node is excited, the low capacitance of the sheath

causes the axonal potential to “jump” from one node of Ranvier to the next, a process known as saltatory conduction. Owing to the low capacitance of the sheath, little energy is required for the remaining membrane between depolarized nodes, which leads to the transmission of action potentials (Seidl, 2014). The propagation of action potentials depends on rapid depolarization and repolarization of the axolemma; these actions are performed by Na⁺ and K⁺ channels that aggregate at nodes. Na⁺ channels act as pore-forming protein subunits that regulate ion flux across membranes and, together with K⁺ channels, regulate neuronal excitability. In addition to ion channels, Ranvier nodes are rich in cytoskeletal and scaffolding proteins such as ankyrin G (ankG) and β IV spectrin, which together link Na⁺ and K⁺ channels to underlying cytoskeletons. These scaffolds also connect ion channels to a diverse and rich extracellular matrix (ECM) through CAMs (the 186-kDa isoform of neurofascin (NF186) and NrCAM) (Rasband and Peles, 2016). However, there is still no consensus on the specific assembly process of Ranvier nodes in the CNS. Three potential mechanisms have been proposed: (1) aggregation of NF186 by ECM derived from glial cells; (2) restriction of nodal protein mobility through paranodal axoglial barriers; and (3) stabilization of Na⁺ channels through axonal chondroitin sulfate (CS). For more information, please refer to the study by Susuki et al. (2013).

4 Another driving force for myelin extension—myelin compaction

However, the molecular mechanism that drives lateral extension of the inner tongue is not well understood. One hypothesis is that actin polymerization at the leading edge drives the process of wrapping, whereas actin depolymerization reduces surface tension, thereby promoting membrane diffusion (Nawaz et al., 2015). Actin polymerization provides power for OL process extension in an Arp2/3-dependent manner while actin depolymerization drives myelin phospholipid wrapping (Zuchero et al., 2015).

In addition to the aforementioned possibilities, we propose a novel hypothesis that myelin compaction provides the driving force for the expansion of myelin. To illustrate this concept, an elastic plastic

container should be filled with water and pressure should be applied to one side. The water will be forced to spread in the opposite direction due to pressure, leading to the expansion of the container. The lateral expansion of myelin is similar. Owing to the “zipper” effect of MBP, excess cytoplasmic and membrane-binding proteins are restricted from entering the dense myelin layer and therefore are squeezed into the uncompacted area (Raasakka et al., 2017). Owing to the increase in the uncompacted area content, the myelin membrane is forced to expand and extend in other directions (Fig. 4). That is, the force generated by the polymerization of MBP is used to propel the myelin membrane forward, similar to how actin polymerization drives the leading edge in migratory cells (Bakhti et al., 2014).




Fig. 4 Force generated by the polymerization of myelin basic protein (MBP) to propel the myelin membrane forward.

The high lateral mobility of myelin lipids makes this hypothesis feasible. Strictly speaking, myelin lipids reduce the adhesiveness of the inner tongue as it crawls under the formed myelin. Notably, myelin has a very high lipid content when compared with other plasma membranes, accounting for 70%–75% of its dry weight, and an unusual lipid composition with a molar ratio of approximately 2:2:1:1 for cholesterol, phospholipids, galactolipids, and plasmalogens (Norton and Poduslo, 1973). Ceramide galactosyltransferase-deficient mice do not synthesize galactocerebroside or sulfatide, but microscopic and morphometric analyses revealed slightly thinner sheaths in the ventral region of the spinal cord although the myelin had a normal ultrastructural appearance (Coetzee et al., 1996). However, cholesterol seems to be a rate-limiting factor in myelin formation, as demonstrated in mutated mice and zebrafish lacking an essential enzyme for cholesterol biosynthesis (Saher et al., 2005; Mathews et al., 2014). Squalene synthase activity is the first step in synthesizing cholesterol (a major lipid component essential for myelin fluidity). OL-specific deletion of squalene synthase results in a significant

decrease in the myelin formation rate (Saher et al., 2005).

Furthermore, as mentioned above, shiverer mice, which lack MBP expression, form less myelin and less stable myelin than wild-type mice. The myelin formed by shiverer mice consists of loosely compacted layers and is at most four layers thick in early development (Roach et al., 1985; Snaidero et al., 2014). Although some basic structural elements of the myelin remain, such as RCs and axon–glial connections, most of the axonal surface is completely devoid of myelin (Rosenbluth, 1980). This is very different from the 160 layers found under normal conditions. This implies that the absence of MBP does not affect the structural integrity of the myelin but rather results in a loss of extension power, leading to a decrease in the number of layers. Strangely, on Day 21, there was no statistically significant difference in the number of myelinated axons with non-compacted layers between optic nerves of wild-type and shiverer heterozygote (Shiv \pm) mice, although Shiv \pm mice had more (Snaidero et al., 2014). This indicates that MBP regulates the speed of compaction in early development, while in the later stages, the speed may be driven by hydrostatic pressure built up by MBP (Stadelmann et al., 2019).

5 Conclusions

Despite advancements in modern technology, our understanding of myelin development in the CNS remains limited. In the human CNS, myelin proteomics has revealed the presence of over 800 proteins (Gargareta et al., 2022), whereas in zebrafish, a significantly higher number (over 1000) of proteins have been identified (Siems et al., 2021); however, current research is focused mainly on high-content proteins. There has been little research on trace proteins, and their roles in myelin formation, growth, functional maintenance, and repair remain unknown. Determining whether these trace proteins play a decisive role in myelin formation and growth poses new challenges for future research. In addition to its structure and function, the mechanism of myelin injury remains to be further studied. Neuroinflammation has been shown to cause changes in myelin. OLs, astrocytes, and microglia are the main glial cells in the CNS. The crosstalk between these cells and their

effects on the formation and destruction of myelin are still unknown. Although we still face important challenges, as the mechanism of myelin formation becomes clearer, the roles of myelin in CNS diseases and even mental disorders may be revealed. This will help the development of new treatments for related diseases.

Acknowledgments

This research was supported by the National Natural Science Foundation of China (No. U21A20400), the Natural Science Foundation of Beijing (No. 8217153264), and the Key Project of Beijing University of Chinese Medicine (No. 2022-JYB-JBZR-004), China.

Author contributions

Writing – original draft: Qi SHAO and Simin CHEN. Visualization: Qi SHAO, Yuyu SHI, and Zijin SUN. Conceptualization: Qingguo WANG and Tian XU. Writing – review and editing: Fafeng CHENG and Xueqian WANG. Funding acquisition: Qingguo WANG and Xueqian WANG. All authors have read and agreed to the published version of the manuscript.

Compliance with ethics guidelines

Qi SHAO, Simin CHEN, Tian XU, Yuyu SHI, Zijin SUN, Qingguo WANG, Xueqian WANG, and Fafeng CHENG declare that they have no conflicts of interest.

This article does not contain any studies with human or animal subjects performed by any of the authors.

References

- Aggarwal S, Yurlova L, Simons M, 2011. Central nervous system myelin: structure, synthesis and assembly. *Trends Cell Biol*, 21(10):585-593. <https://doi.org/10.1016/j.tcb.2011.06.004>
- Bakhti M, Snaidero N, Schneider D, et al., 2013. Loss of electrostatic cell-surface repulsion mediates myelin membrane adhesion and compaction in the central nervous system. *Proc Natl Acad Sci USA*, 110(8):3143-3148. <https://doi.org/10.1073/pnas.1220104110>
- Bakhti M, Aggarwal S, Simons M, 2014. Myelin architecture: zipper membranes tightly together. *Cell Mol Life Sci*, 71(7):1265-1277. <https://doi.org/10.1007/s00018-013-1492-0>
- Barateiro A, Brites D, Fernandes A, 2016. Oligodendrocyte development and myelination in neurodevelopment: molecular mechanisms in health and disease. *Curr Pharm Des*, 22(6):656-679. <https://doi.org/10.2174/1381612822666151204000636>
- Bechler ME, Byrne L, Ffrench-Constant C, 2015. CNS myelin sheath lengths are an intrinsic property of oligodendrocytes. *Curr Biol*, 25(18):2411-2416. <https://doi.org/10.1016/j.cub.2015.07.056>
- Bifulco M, Laezza C, Stingo S, et al., 2002. 2',3'-Cyclic nucleotide 3'-phosphodiesterase: a membrane-bound, microtubule-associated protein and membrane anchor for tubulin. *Proc Natl Acad Sci USA*, 99(4):1807-1812. <https://doi.org/10.1073/pnas.042678799>
- Brady ST, Witt AS, Kirkpatrick LL, et al., 1999. Formation of compact myelin is required for maturation of the axonal cytoskeleton. *J Neurosci*, 19(17):7278-7288. <https://doi.org/10.1523/JNEUROSCI.19-17-07278.1999>
- Campagnoni AT, 1988. Molecular biology of myelin proteins from the central nervous system. *J Neurochem*, 51(1):1-14. <https://doi.org/10.1111/j.1471-4159.1988.tb04827.x>
- Chow E, Mottahedeh J, Prins M, et al., 2005. Disrupted compaction of CNS myelin in an OSP/Claudin-11 and PLP/DM20 double knockout mouse. *Mol Cell Neurosci*, 29(3):405-413. <https://doi.org/10.1016/j.mcn.2005.03.007>
- Coetzee T, Fujita N, Dupree J, et al., 1996. Myelination in the absence of galactocerebroside and sulfatide: normal structure with abnormal function and regional instability. *Cell*, 86(2):209-219. [https://doi.org/10.1016/s0092-8674\(00\)80093-8](https://doi.org/10.1016/s0092-8674(00)80093-8)
- Correale J, Ysrraelit MC, 2022. Multiple sclerosis and aging: the dynamics of demyelination and remyelination. *ASN Neuro*, 14:17590914221118502. <https://doi.org/10.1177/17590914221118502>
- de Faria O, Pivonkova H, Varga B, et al., 2021. Periods of synchronized myelin changes shape brain function and plasticity. *Nat Neurosci*, 24(11):1508-1521. <https://doi.org/10.1038/s41593-021-00917-2>
- Denninger AR, Breglio A, Maheras KJ, et al., 2015. Claudin-11 tight junctions in myelin are a barrier to diffusion and lack strong adhesive properties. *Biophys J*, 109(7):1387-1397. <https://doi.org/10.1016/j.bpj.2015.08.012>
- Devaux J, Gow A, 2008. Tight junctions potentiate the insulative properties of small CNS myelinated axons. *J Cell Biol*, 183(5):909-921. <https://doi.org/10.1083/jcb.200808034>
- Edgar JM, McLaughlin M, Werner HB, et al., 2009. Early ultrastructural defects of axons and axon–glia junctions in mice lacking expression of *Cnp1*. *Glia*, 57(16):1815-1824. <https://doi.org/10.1002/glia.20893>
- Edgar JM, McGowan E, Chapple KJ, et al., 2021. Río-Hortega's drawings revisited with fluorescent protein defines a cytoplasm-filled channel system of CNS myelin. *J Anat*, 239(6):1241-1255. <https://doi.org/10.1111/joa.13577>
- Fehmi J, Scherer SS, Willison HJ, et al., 2018. Nodes, paranodes and neuropathies. *J Neurol Neurosurg Psychiatry*, 89(1):61-71. <https://doi.org/10.1136/jnnp-2016-315480>
- Gargareta VI, Reuschenbach J, Siems SB, et al., 2022. Conservation and divergence of myelin proteome and oligodendrocyte transcriptome profiles between humans and mice. *eLife*, 11:e77019.

- <https://doi.org/10.7554/eLife.77019>
- Geoffroy CG, Zheng BH, 2014. Myelin-associated inhibitors in axonal growth after CNS injury. *Curr Opin Neurobiol*, 27:31-38.
<https://doi.org/10.1016/j.conb.2014.02.012>
- Gotoh L, Inoue K, Helman G, et al., 2014. *GJC2* promoter mutations causing Pelizaeus-Merzbacher-like disease. *Mol Genet Metab*, 111(3):393-398.
<https://doi.org/10.1016/j.ymgme.2013.12.001>
- Gow A, Southwood CM, Li JS, et al., 1999. CNS myelin and Sertoli cell tight junction strands are absent in *Osp/claudin-11* null mice. *Cell*, 99(6):649-659.
[https://doi.org/10.1016/s0092-8674\(00\)81553-6](https://doi.org/10.1016/s0092-8674(00)81553-6)
- Gravel M, Peterson J, Yong VW, et al., 1996. Overexpression of 2',3'-cyclic nucleotide 3'-phosphodiesterase in transgenic mice alters oligodendrocyte development and produces aberrant myelination. *Mol Cell Neurosci*, 7(6):453-466.
<https://doi.org/10.1006/mcne.1996.0033>
- Gravel M, Robert F, Kottis V, et al., 2009. 2',3'-Cyclic nucleotide 3'-phosphodiesterase: a novel RNA-binding protein that inhibits protein synthesis. *J Neurosci Res*, 87(5):1069-1079.
<https://doi.org/10.1002/jnr.21939>
- Harauz G, Ishiyama N, Hill CMD, et al., 2004. Myelin basic protein-diverse conformational states of an intrinsically unstructured protein and its roles in myelin assembly and multiple sclerosis. *Micron*, 35(7):503-542.
<https://doi.org/10.1016/j.micron.2004.04.005>
- Harry GJ, Toews AD, 1998. Myelination, dysmyelination, and demyelination. In: Slikker W Jr, Chang LW (Eds.), *Handbook of Developmental Neurotoxicology*. Academic Press, San Diego, p.87-115.
<https://doi.org/10.1016/B978-012648860-9.50007-8>
- Hines JH, Ravanelli AM, Schwindt R, et al., 2015. Neuronal activity biases axon selection for myelination *in vivo*. *Nat Neurosci*, 18(5):683-689.
<https://doi.org/10.1038/nn.3992>
- Hou AN, Qu SS, Fu JH, 2020. Expression of connexin 47 and its effects on the myelination of oligodendrocytes in hypoxia-ischemia-induced periventricular leukomalacia in newborn rats. *J China Med Univ*, 49(4):306-312 (in Chinese).
<https://doi.org/10.12007/j.issn.0258-4646.2020.04.004>
- Hughes EG, Kang SH, Fukaya M, et al., 2013. Oligodendrocyte progenitors balance growth with self-repulsion to achieve homeostasis in the adult brain. *Nat Neurosci*, 16(6):668-676.
<https://doi.org/10.1038/nn.3390>
- Kagawa T, Ikenaka K, Inoue Y, et al., 1994. Glial cell degeneration and hypomyelination caused by overexpression of myelin proteolipid protein gene. *Neuron*, 13(2):427-442.
[https://doi.org/10.1016/0896-6273\(94\)90358-1](https://doi.org/10.1016/0896-6273(94)90358-1)
- Kang M, Yao Y, 2022. Laminin regulates oligodendrocyte development and myelination. *Glia*, 70(3):414-429.
<https://doi.org/10.1002/glia.24117>
- Karthigasan J, Kirschner DA, 1988. Membrane interactions are altered in myelin isolated from central and peripheral nervous system tissues. *J Neurochem*, 51(1):228-236.
<https://doi.org/10.1111/j.1471-4159.1988.tb04860.x>
- Kerman BE, Kim HJ, Padmanabhan K, et al., 2015. *In vitro* myelin formation using embryonic stem cells. *Development*, 142(12):2213-2225.
<https://doi.org/10.1242/dev.116517>
- Kippert A, Trajkovic K, Rajendran L, et al., 2007. Rho regulates membrane transport in the endocytic pathway to control plasma membrane specialization in oligodendroglial cells. *J Neurosci*, 27(13):3560-3570.
<https://doi.org/10.1523/JNEUROSCI.4926-06.2007>
- Kirschner DA, Inouye H, Ganser AL, et al., 1989. Myelin membrane structure and composition correlated: a phylogenetic study. *J Neurochem*, 53(5):1599-1609.
<https://doi.org/10.1111/j.1471-4159.1989.tb08558.x>
- Kister A, Kister I, 2023. Overview of myelin, major myelin lipids, and myelin-associated proteins. *Front Chem*, 10:1041961.
<https://doi.org/10.3389/fchem.2022.1041961>
- Klugmann M, Schwab MH, Pühlhofer A, et al., 1997. Assembly of CNS myelin in the absence of proteolipid protein. *Neuron*, 18(1):59-70.
[https://doi.org/10.1016/s0896-6273\(01\)80046-5](https://doi.org/10.1016/s0896-6273(01)80046-5)
- Krafft PR, Bailey EL, Lekic T, et al., 2012. Etiology of stroke and choice of models. *Int J Stroke*, 7(5):398-406.
<https://doi.org/10.1111/j.1747-4949.2012.00838.x>
- Kuhn S, Gritti L, Crooks D, et al., 2019. Oligodendrocytes in development, myelin generation and beyond. *Cells*, 8(11):1424.
<https://doi.org/10.3390/cells8111424>
- Lapato AS, Tiwari-Woodruff SK, 2018. Connexins and pannexins: at the junction of neuro-glial homeostasis & disease. *J Neurosci Res*, 96(1):31-44.
<https://doi.org/10.1002/jnr.24088>
- Lappe-Siefke C, Goebbels S, Gravel M, et al., 2003. Disruption of *Cnp1* uncouples oligodendroglial functions in axonal support and myelination. *Nat Genet*, 33(3):366-374.
<https://doi.org/10.1038/ng1095>
- Laursen LS, Chan CW, Ffrench-Constant C, 2011. Translation of myelin basic protein mRNA in oligodendrocytes is regulated by integrin activation and hnRNP-K. *J Cell Biol*, 192(5):797-811.
<https://doi.org/10.1083/jcb.201007014>
- Lee J, Gravel M, Zhang RL, et al., 2005. Process outgrowth in oligodendrocytes is mediated by CNP, a novel microtubule assembly myelin protein. *J Cell Biol*, 170(4):661-673.
<https://doi.org/10.1083/jcb.200411047>
- Lee S, Leach MK, Redmond SA, et al., 2012. A culture system to study oligodendrocyte myelination processes using engineered nanofibers. *Nat Methods*, 9(9):917-922.
<https://doi.org/10.1038/nmeth.2105>
- Lee S, Chong SYC, Tuck SJ, et al., 2013. A rapid and reproducible assay for modeling myelination by oligodendrocytes using engineered nanofibers. *Nat Protoc*, 8(4):771-782.
<https://doi.org/10.1038/nprot.2013.039>

- Lepiemme F, Stoufflet J, Javier-Torrent M, et al., 2022. Oligodendrocyte precursors guide interneuron migration by unidirectional contact repulsion. *Science*, 376(6595):eabn6204. <https://doi.org/10.1126/science.abn6204>
- Liu H, Yang YY, Xia YG, et al., 2017. Aging of cerebral white matter. *Ageing Res Rev*, 34:64-76. <https://doi.org/10.1016/j.arr.2016.11.006>
- Lüders KA, Nessler S, Kusch K, et al., 2019. Maintenance of high proteolipid protein level in adult central nervous system myelin is required to preserve the integrity of myelin and axons. *Glia*, 67(4):634-649. <https://doi.org/10.1002/glia.23549>
- Maglione M, Tress O, Haas B, et al., 2010. Oligodendrocytes in mouse corpus callosum are coupled via gap junction channels formed by connexin47 and connexin32. *Glia*, 58(9):1104-1117. <https://doi.org/10.1002/glia.20991>
- Mathews ES, Mawdsley DJ, Walker M, et al., 2014. Mutation of 3-hydroxy-3-methylglutaryl CoA synthase I reveals requirements for isoprenoid and cholesterol synthesis in oligodendrocyte migration arrest, axon wrapping, and myelin gene expression. *J Neurosci*, 34(9):3402-3412. <https://doi.org/10.1523/JNEUROSCI.4587-13.2014>
- Mayoral SR, Etxeberria A, Shen YAA, et al., 2018. Initiation of CNS myelination in the optic nerve is dependent on axon caliber. *Cell Rep*, 25(3):544-550.e3. <https://doi.org/10.1016/j.celrep.2018.09.052>
- McKerracher L, Rosen KM, 2015. MAG, myelin and overcoming growth inhibition in the CNS. *Front Mol Neurosci*, 8:51. <https://doi.org/10.3389/fnmol.2015.00051>
- McNamara NB, Munro DAD, Bestard-Cuche N, et al., 2023. Microglia regulate central nervous system myelin growth and integrity. *Nature*, 613(7942):120-129. <https://doi.org/10.1038/s41586-022-05534-y>
- Min Y, Kristiansen K, Boggs JM, et al., 2009. Interaction forces and adhesion of supported myelin lipid bilayers modulated by myelin basic protein. *Proc Natl Acad Sci USA*, 106(9):3154-3159. <https://doi.org/10.1073/pnas.0813110106>
- Moscarello MA, 1997. Myelin basic protein, the “executive” molecule of the myelin membrane. In: Juurlink BHJ, Devon RM, Doucette JR, et al. (Eds.), *Verge Cell Biology and Pathology of Myelin: Evolving Biological Concepts and Therapeutic Approaches*. Springer, Boston, p.13-25. https://doi.org/10.1007/978-1-4615-5949-8_2
- Müller C, Bauer NM, Schäfer I, et al., 2013. Making myelin basic protein—from mRNA transport to localized translation. *Front Cell Neurosci*, 7:169. <https://doi.org/10.3389/fncel.2013.00169>
- Nave KA, 2010. Myelination and support of axonal integrity by glia. *Nature*, 468(7321):244-252. <https://doi.org/10.1038/nature09614>
- Nawaz S, Sánchez P, Schmitt S, et al., 2015. Actin filament turnover drives leading edge growth during myelin sheath formation in the central nervous system. *Dev Cell*, 34(2):139-151. <https://doi.org/10.1016/j.devcel.2015.05.013>
- Norton WT, Poduslo SE, 1973. Myelination in rat brain: changes in myelin composition during brain maturation. *J Neurochem*, 21(4):759-773. <https://doi.org/10.1111/j.1471-4159.1973.tb07520.x>
- Pajevic S, Plenz D, Basser PJ, et al., 2023. Oligodendrocyte-mediated myelin plasticity and its role in neural synchronization. *eLife*, 12:e81982. <https://doi.org/10.7554/eLife.81982>
- Pedraza L, Huang JK, Colman D, 2009. Disposition of axonal caspr with respect to glial cell membranes: implications for the process of myelination. *J Neurosci Res*, 87(15):3480-3491. <https://doi.org/10.1002/jnr.22004>
- Pfeiffer SE, Warrington AE, Bansal R, 1993. The oligodendrocyte and its many cellular processes. *Trends Cell Biol*, 3(6):191-197. [https://doi.org/10.1016/0962-8924\(93\)90213-k](https://doi.org/10.1016/0962-8924(93)90213-k)
- Pillai AM, Thaxton C, Pribisko AL, et al., 2009. Spatiotemporal ablation of myelinating glia-specific *neurofascin* (*Nfasc*^{NF155}) in mice reveals gradual loss of paranodal axoglial junctions and concomitant disorganization of axonal domains. *J Neurosci Res*, 87(8):1773-1793. <https://doi.org/10.1002/jnr.22015>
- Pinatel D, Faivre-Sarrailh C, 2021. Assembly and function of the juxtaparanodal Kv1 complex in health and disease. *Life*, 11(1):8. <https://doi.org/10.3390/life11010008>
- Raasakka A, Ruskamo S, Kowal J, et al., 2017. Membrane association landscape of myelin basic protein portrays formation of the myelin major dense line. *Sci Rep*, 7:4974. <https://doi.org/10.1038/s41598-017-05364-3>
- Raasakka A, Ruskamo S, Kowal J, et al., 2019. Molecular structure and function of myelin protein P0 in membrane stacking. *Sci Rep*, 9:642. <https://doi.org/10.1038/s41598-018-37009-4>
- Rasband MN, Peles E, 2016. The nodes of Ranvier: molecular assembly and maintenance. *Cold Spring Harb Perspect Biol*, 8(3):a020495. <https://doi.org/10.1101/cshperspect.a020495>
- Roach A, Takahashi N, Pravtcheva D, et al., 1985. Chromosomal mapping of mouse myelin basic protein gene and structure and transcription of the partially deleted gene in shiverer mutant mice. *Cell*, 42(1):149-155. [https://doi.org/10.1016/s0092-8674\(85\)80110-0](https://doi.org/10.1016/s0092-8674(85)80110-0)
- Rosenberg SS, Kelland EE, Tokar E, et al., 2008. The geometric and spatial constraints of the microenvironment induce oligodendrocyte differentiation. *Proc Natl Acad Sci USA*, 105(38):14662-14667. <https://doi.org/10.1073/pnas.0805640105>
- Rosenbluth J, 1980. Central myelin in the mouse mutant shiverer. *J Comp Neurol*, 194(3):639-648. <https://doi.org/10.1002/cne.901940310>
- Rosenbluth J, Nave KA, Mierzwa A, et al., 2006. Subtle myelin defects in PLP-null mice. *Glia*, 54(3):172-182. <https://doi.org/10.1002/glia.20370>

- Ruskamo S, Raasakka A, Pedersen JS, et al., 2022. Human myelin proteolipid protein structure and lipid bilayer stacking. *Cell Mol Life Sci*, 79(8):419.
<https://doi.org/10.1007/s00018-022-04428-6>
- Saher G, Brügger B, Lappe-Siefke C, et al., 2005. High cholesterol level is essential for myelin membrane growth. *Nat Neurosci*, 8(4):468-475.
<https://doi.org/10.1038/nn1426>
- Salzer JL, Zalc B, 2016. Myelination. *Curr Biol*, 26(20):R971-R975.
<https://doi.org/10.1016/j.cub.2016.07.074>
- Sargiannidou I, Kim GH, Kyriakoudi S, et al., 2015. A start codon CMT1X mutation associated with transient encephalomyelitis causes complete loss of Cx32. *Neurogenetics*, 16(3):193-200.
<https://doi.org/10.1007/s10048-015-0442-4>
- Seidl AH, 2014. Regulation of conduction time along axons. *Neuroscience*, 276:126-134.
<https://doi.org/10.1016/j.neuroscience.2013.06.047>
- Sherman DL, Tait S, Melrose S, et al., 2005. Neurofascins are required to establish axonal domains for saltatory conduction. *Neuron*, 48(5):737-742.
<https://doi.org/10.1016/j.neuron.2005.10.019>
- Siems SB, Jahn O, Hoodless LJ, et al., 2021. Proteome profile of myelin in the zebrafish brain. *Front Cell Dev Biol*, 9:640169.
<https://doi.org/10.3389/fcell.2021.640169>
- Simons M, Nave KA, 2016. Oligodendrocytes: myelination and axonal support. *Cold Spring Harb Perspect Biol*, 8(1):a020479.
<https://doi.org/10.1101/cshperspect.a020479>
- Smirnova EV, Rakitina TV, Ziganshin RH, et al., 2021. Comprehensive atlas of the myelin basic protein interaction landscape. *Biomolecules*, 11(11):1628.
<https://doi.org/10.3390/biom11111628>
- Snaidero N, Möbius W, Czopka T, et al., 2014. Myelin membrane wrapping of CNS axons by PI(3,4,5)P3-dependent polarized growth at the inner tongue. *Cell*, 156(1-2):277-290.
<https://doi.org/10.1016/j.cell.2013.11.044>
- Snaidero N, Velte C, Myllykoski M, et al., 2017. Antagonistic functions of MBP and CNP establish cytosolic channels in CNS myelin. *Cell Rep*, 18(2):314-323.
<https://doi.org/10.1016/j.celrep.2016.12.053>
- Sobottka B, Ziegler U, Kaech A, et al., 2011. CNS live imaging reveals a new mechanism of myelination: the liquid croissant model. *Glia*, 59(12):1841-1849.
<https://doi.org/10.1002/glia.21228>
- Stadelmann C, Timmler S, Barrantes-Freer A, et al., 2019. Myelin in the central nervous system: structure, function, and pathology. *Physiol Rev*, 99(3):1381-1431.
<https://doi.org/10.1152/physrev.00031.2018>
- Susuki K, Chang KJ, Zollinger DR, et al., 2013. Three mechanisms assemble central nervous system nodes of Ranvier. *Neuron*, 78(3):469-482.
<https://doi.org/10.1016/j.neuron.2013.03.005>
- Susuki K, Otani Y, Rasband MN, 2016. Submembranous cytoskeletons stabilize nodes of Ranvier. *Exp Neurol*, 283:446-451.
<https://doi.org/10.1016/j.expneurol.2015.11.012>
- Tepavčević V, Lubetzki C, 2022. Oligodendrocyte progenitor cell recruitment and remyelination in multiple sclerosis: the more, the merrier? *Brain*, 145(12):4178-4192.
<https://doi.org/10.1093/brain/awac307>
- Thomason EJ, Escalante M, Osterhout DJ, et al., 2020. The oligodendrocyte growth cone and its actin cytoskeleton: a fundamental element for progenitor cell migration and CNS myelination. *Glia*, 68(7):1329-1346.
<https://doi.org/10.1002/glia.23735>
- Trapp BD, 1990. Myelin-associated glycoprotein location and potential functions. *Ann N Y Acad Sci*, 605(1):29-43.
<https://doi.org/10.1111/j.1749-6632.1990.tb42378.x>
- Tress O, Maglione M, Zlomuzica A, et al., 2011. Pathologic and phenotypic alterations in a mouse expressing a connexin47 missense mutation that causes Pelizaeus-Merzbacher-like disease in humans. *PLoS Genet*, 7(7):e1002146.
<https://doi.org/10.1371/journal.pgen.1002146>
- Wang SQ, Wang YX, Zou SQ, 2022. A glance at the molecules that regulate oligodendrocyte myelination. *Curr Issues Mol Biol*, 44(5):2194-2216.
<https://doi.org/10.3390/cimb44050149>
- Waxman SG, Bennett MVL, 1972. Relative conduction velocities of small myelinated and non-myelinated fibres in the central nervous system. *Nat New Biol*, 238(85):217-219.
<https://doi.org/10.1038/newbio238217a0>
- Weimbs T, Stoffel W, 1992. Proteolipid protein (PLP) of CNS myelin: positions of free, disulfide-bonded, and fatty acid thioester-linked cysteine residues and implications for the membrane topology of PLP. *Biochemistry*, 31(49):12289-12296.
<https://doi.org/10.1021/bi00164a002>
- Zalc B, Goujet D, Colman D, 2008. The origin of the myelination program in vertebrates. *Curr Biol*, 18(12):R511-R512.
<https://doi.org/10.1016/j.cub.2008.04.010>
- Zuchero JB, Fu MM, Sloan SA, et al., 2015. CNS myelin wrapping is driven by actin disassembly. *Dev Cell*, 34(2):152-167.
<https://doi.org/10.1016/j.devcel.2015.06.011>